

Raising to generic powers

Jonathan Kirby

University of Oxford

Modnet conference, Barcelona, 2008

Abstract

We prove unconditionally a Schanuel property for raising to a generic real power, leading to the hope that the real field with a generic power function can be proved to be decidable. This is joint work with A.J. Wilkie and Martin Bays.

Outline

- 1 Motivation – Decidability
- 2 Schanuel Properties
- 3 Proofs

Outline

1 Motivation – Decidability

2 Schanuel Properties

3 Proofs

Decidability of \mathbb{R}

Theorem (Tarski 1949)

The theory of the real field $\langle \mathbb{R}; +, \cdot \rangle$ is decidable.

Proof uses:

- model completeness
- A decision procedure for \exists -sentences

Tarski asked: is $\mathbb{R}_{\text{exp}} = \langle \mathbb{R}; +, \cdot, \text{exp} \rangle$ decidable?

Decidability of \mathbb{R}

Theorem (Tarski 1949)

The theory of the real field $\langle \mathbb{R}; +, \cdot \rangle$ is decidable.

Proof uses:

- model completeness
- A decision procedure for \exists -sentences

Tarski asked: is $\mathbb{R}_{\text{exp}} = \langle \mathbb{R}; +, \cdot, \text{exp} \rangle$ decidable?

Decidability of \mathbb{R}

Theorem (Tarski 1949)

The theory of the real field $\langle \mathbb{R}; +, \cdot \rangle$ is decidable.

Proof uses:

- model completeness
- A decision procedure for \exists -sentences

Tarski asked: is $\mathbb{R}_{\text{exp}} = \langle \mathbb{R}; +, \cdot, \text{exp} \rangle$ decidable?

Decidability of \mathbb{R}

Theorem (Tarski 1949)

The theory of the real field $\langle \mathbb{R}; +, \cdot \rangle$ is decidable.

Proof uses:

- model completeness
- A decision procedure for \exists -sentences

Tarski asked: is $\mathbb{R}_{\text{exp}} = \langle \mathbb{R}; +, \cdot, \text{exp} \rangle$ decidable?

Decidability of \mathbb{R}

Theorem (Tarski 1949)

The theory of the real field $\langle \mathbb{R}; +, \cdot \rangle$ is decidable.

Proof uses:

- model completeness
- A decision procedure for \exists -sentences

Tarski asked: is $\mathbb{R}_{\text{exp}} = \langle \mathbb{R}; +, \cdot, \text{exp} \rangle$ decidable?

Decidability of \mathbb{R}_{exp}

Theorem (Wilkie 1996)

\mathbb{R}_{exp} is model-complete and o-minimal

Theorem (Macintyre, Wilkie 1996)

Assuming Schanuel's Conjecture, there is a decision procedure for \exists -sentences.

Corollary

Conditionally, \mathbb{R}_{exp} is decidable.

Decidability of \mathbb{R}_{exp}

Theorem (Wilkie 1996)

\mathbb{R}_{exp} is model-complete and o-minimal

Theorem (Macintyre, Wilkie 1996)

Assuming Schanuel's Conjecture, there is a decision procedure for \exists -sentences.

Corollary

Conditionally, \mathbb{R}_{exp} is decidable.

Decidability of \mathbb{R}_{exp}

Theorem (Wilkie 1996)

\mathbb{R}_{exp} is model-complete and o-minimal

Theorem (Macintyre, Wilkie 1996)

Assuming Schanuel's Conjecture, there is a decision procedure for \exists -sentences.

Corollary

Conditionally, \mathbb{R}_{exp} is decidable.

Decidability of other functions

Question (Jones, et al.)

Can we unconditionally prove decidability for $\langle \mathbb{R}; +, \cdot, f \rangle$ for *some* (interesting) analytic function f ?

Raising to a power $\lambda \in \mathbb{R}$

For $y > 0$, $y^\lambda = \exp(\lambda \log y)$

$\mathbb{R}_\lambda = \langle \mathbb{R}; +, \cdot, (-)^\lambda \rangle$ is o-minimal and model complete (Wilkie / Miller)

Work in progress – Jones, Servi

- Schanuel Property for $(-)^{\lambda} \implies$ decidability of \mathbb{R}_λ
- (if λ is a recursive real – otherwise decidability modulo λ)

Decidability of other functions

Question (Jones, et al.)

Can we unconditionally prove decidability for $\langle \mathbb{R}; +, \cdot, f \rangle$ for *some* (interesting) analytic function f ?

Raising to a power $\lambda \in \mathbb{R}$

For $y > 0$, $y^\lambda = \exp(\lambda \log y)$

$\mathbb{R}_\lambda = \langle \mathbb{R}; +, \cdot, (-)^\lambda \rangle$ is o-minimal and model complete (Wilkie / Miller)

Work in progress – Jones, Servi

- Schanuel Property for $(-)^{\lambda} \implies$ decidability of \mathbb{R}_λ
- (if λ is a recursive real – otherwise decidability modulo λ)

Decidability of other functions

Question (Jones, et al.)

Can we unconditionally prove decidability for $\langle \mathbb{R}; +, \cdot, f \rangle$ for *some* (interesting) analytic function f ?

Raising to a power $\lambda \in \mathbb{R}$

For $y > 0$, $y^\lambda = \exp(\lambda \log y)$

$\mathbb{R}_\lambda = \langle \mathbb{R}; +, \cdot, (-)^\lambda \rangle$ is o-minimal and model complete (Wilkie / Miller)

Work in progress – Jones, Servi

- Schanuel Property for $(-)^{\lambda} \implies$ decidability of \mathbb{R}_λ
- (if λ is a recursive real – otherwise decidability modulo λ)

Decidability of other functions

Question (Jones, et al.)

Can we unconditionally prove decidability for $\langle \mathbb{R}; +, \cdot, f \rangle$ for *some* (interesting) analytic function f ?

Raising to a power $\lambda \in \mathbb{R}$

For $y > 0$, $y^\lambda = \exp(\lambda \log y)$

$\mathbb{R}_\lambda = \langle \mathbb{R}; +, \cdot, (-)^\lambda \rangle$ is o-minimal and model complete (Wilkie / Miller)

Work in progress – Jones, Servi

- Schanuel Property for $(-)^{\lambda} \implies$ decidability of \mathbb{R}_λ
- (if λ is a recursive real – otherwise decidability modulo λ)

Decidability of other functions

Question (Jones, et al.)

Can we unconditionally prove decidability for $\langle \mathbb{R}; +, \cdot, f \rangle$ for *some* (interesting) analytic function f ?

Raising to a power $\lambda \in \mathbb{R}$

For $y > 0$, $y^\lambda = \exp(\lambda \log y)$

$\mathbb{R}_\lambda = \langle \mathbb{R}; +, \cdot, (-)^\lambda \rangle$ is o-minimal and model complete (Wilkie / Miller)

Work in progress – Jones, Servi

- Schanuel Property for $(-)^{\lambda} \implies$ decidability of \mathbb{R}_λ
- (if λ is a recursive real – otherwise decidability modulo λ)

Outline

1 Motivation – Decidability

2 Schanuel Properties

3 Proofs

Schanuel Property for raising to a power

Theorem (Bays, Kirby, Wilkie)

Let $\lambda \in \mathbb{R}$ be exponentially transcendental, let $\bar{y} \in (\mathbb{R}_{>0})^n$, and suppose \bar{y} is multiplicatively independent. Then

$$\text{td } \mathbb{Q}(\bar{y}, \bar{y}^\lambda, \lambda) / \mathbb{Q}(\lambda) \geq n.$$

- λ is exponentially transcendental iff it does not lie in the prime model of \mathbb{R}_{exp} .
- Co-countably many reals are exponentially transcendental.
- No known exponentially transcendental reals!
- Cantor's argument gives a recursive exponentially transcendental real.

Schanuel Property for raising to a power

Theorem (Bays, Kirby, Wilkie)

Let $\lambda \in \mathbb{R}$ be exponentially transcendental, let $\bar{y} \in (\mathbb{R}_{>0})^n$, and suppose \bar{y} is multiplicatively independent. Then

$$\text{td } \mathbb{Q}(\bar{y}, \bar{y}^\lambda, \lambda) / \mathbb{Q}(\lambda) \geq n.$$

- λ is exponentially transcendental iff it does not lie in the prime model of \mathbb{R}_{exp} .
- Co-countably many reals are exponentially transcendental.
- No known exponentially transcendental reals!
- Cantor's argument gives a recursive exponentially transcendental real.

Schanuel Property for raising to a power

Theorem (Bays, Kirby, Wilkie)

Let $\lambda \in \mathbb{R}$ be exponentially transcendental, let $\bar{y} \in (\mathbb{R}_{>0})^n$, and suppose \bar{y} is multiplicatively independent. Then

$$\text{td } \mathbb{Q}(\bar{y}, \bar{y}^\lambda, \lambda) / \mathbb{Q}(\lambda) \geq n.$$

- λ is exponentially transcendental iff it does not lie in the prime model of \mathbb{R}_{exp} .
- Co-countably many reals are exponentially transcendental.
- No known exponentially transcendental reals!
- Cantor's argument gives a recursive exponentially transcendental real.

Schanuel Property for raising to a power

Theorem (Bays, Kirby, Wilkie)

Let $\lambda \in \mathbb{R}$ be exponentially transcendental, let $\bar{y} \in (\mathbb{R}_{>0})^n$, and suppose \bar{y} is multiplicatively independent. Then

$$\text{td } \mathbb{Q}(\bar{y}, \bar{y}^\lambda, \lambda) / \mathbb{Q}(\lambda) \geq n.$$

- λ is exponentially transcendental iff it does not lie in the prime model of \mathbb{R}_{exp} .
- Co-countably many reals are exponentially transcendental.
- No known exponentially transcendental reals!
- Cantor's argument gives a recursive exponentially transcendental real.

Schanuel Property for raising to a power

Theorem (Bays, Kirby, Wilkie)

Let $\lambda \in \mathbb{R}$ be exponentially transcendental, let $\bar{y} \in (\mathbb{R}_{>0})^n$, and suppose \bar{y} is multiplicatively independent. Then

$$\text{td } \mathbb{Q}(\bar{y}, \bar{y}^\lambda, \lambda) / \mathbb{Q}(\lambda) \geq n.$$

- λ is exponentially transcendental iff it does not lie in the prime model of \mathbb{R}_{exp} .
- Co-countably many reals are exponentially transcendental.
- No known exponentially transcendental reals!
- Cantor's argument gives a recursive exponentially transcendental real.

Exponential Transcendence - general definition

Let $\langle F; +, \cdot, \exp \rangle$ be any exponential field.

Definition

$x \in F$ is **exponentially algebraic** in F iff for some $n \in \mathbb{N}$ there are:

- $\bar{x} = (x_1, \dots, x_n) \in F^n$
- $f_1, \dots, f_n \in \mathbb{Z}[\bar{X}, e^{\bar{X}}]$

such that

- $x = x_1$
- $f_i(\bar{x}, e^{\bar{x}}) = 0$ for each $i = 1, \dots, n$

- $$\begin{vmatrix} \frac{\partial f_1}{\partial X_1} & \cdots & \frac{\partial f_1}{\partial X_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial X_1} & \cdots & \frac{\partial f_n}{\partial X_n} \end{vmatrix}(\bar{x}) \neq 0$$

Exponentially Transcendental in $F \iff$ not exponentially algebraic in F

Exponential Transcendence - general definition

Let $\langle F; +, \cdot, \exp \rangle$ be any exponential field.

Definition

$x \in F$ is **exponentially algebraic** in F iff for some $n \in \mathbb{N}$ there are:

- $\bar{x} = (x_1, \dots, x_n) \in F^n$
- $f_1, \dots, f_n \in \mathbb{Z}[\bar{X}, e^{\bar{X}}]$

such that

- $x = x_1$
- $f_i(\bar{x}, e^{\bar{x}}) = 0$ for each $i = 1, \dots, n$

- $$\begin{vmatrix} \frac{\partial f_1}{\partial X_1} & \cdots & \frac{\partial f_1}{\partial X_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial X_1} & \cdots & \frac{\partial f_n}{\partial X_n} \end{vmatrix}(\bar{x}) \neq 0$$

Exponentially Transcendental in $F \iff$ not exponentially algebraic in F

Exponential Transcendence - general definition

Let $\langle F; +, \cdot, \exp \rangle$ be any exponential field.

Definition

$x \in F$ is **exponentially algebraic** in F iff for some $n \in \mathbb{N}$ there are:

- $\bar{x} = (x_1, \dots, x_n) \in F^n$
- $f_1, \dots, f_n \in \mathbb{Z}[\bar{X}, e^{\bar{X}}]$

such that

- $x = x_1$
- $f_i(\bar{x}, e^{\bar{x}}) = 0$ for each $i = 1, \dots, n$

- $$\begin{vmatrix} \frac{\partial f_1}{\partial X_1} & \cdots & \frac{\partial f_1}{\partial X_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial X_1} & \cdots & \frac{\partial f_n}{\partial X_n} \end{vmatrix}(\bar{x}) \neq 0$$

Exponentially Transcendental in $F \iff$ not exponentially algebraic in F

Exponential Transcendence - general definition

Let $\langle F; +, \cdot, \exp \rangle$ be any exponential field.

Definition

$x \in F$ is **exponentially algebraic** in F iff for some $n \in \mathbb{N}$ there are:

- $\bar{x} = (x_1, \dots, x_n) \in F^n$
- $f_1, \dots, f_n \in \mathbb{Z}[\bar{X}, e^{\bar{X}}]$

such that

- $x = x_1$
- $f_i(\bar{x}, e^{\bar{x}}) = 0$ for each $i = 1, \dots, n$

- $$\begin{vmatrix} \frac{\partial f_1}{\partial X_1} & \cdots & \frac{\partial f_1}{\partial X_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial X_1} & \cdots & \frac{\partial f_n}{\partial X_n} \end{vmatrix}(\bar{x}) \neq 0$$

Exponentially Transcendental in $F \iff$ not exponentially algebraic in F

Exponential Transcendence - general definition

Let $\langle F; +, \cdot, \exp \rangle$ be any exponential field.

Definition

$x \in F$ is **exponentially algebraic** in F iff for some $n \in \mathbb{N}$ there are:

- $\bar{x} = (x_1, \dots, x_n) \in F^n$
- $f_1, \dots, f_n \in \mathbb{Z}[\bar{X}, e^{\bar{X}}]$

such that

- $x = x_1$
- $f_i(\bar{x}, e^{\bar{x}}) = 0$ for each $i = 1, \dots, n$

- $$\begin{vmatrix} \frac{\partial f_1}{\partial X_1} & \cdots & \frac{\partial f_1}{\partial X_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial X_1} & \cdots & \frac{\partial f_n}{\partial X_n} \end{vmatrix}(\bar{x}) \neq 0$$

Exponentially Transcendental in $F \iff$ not exponentially algebraic in F

Exponential Transcendence - general definition

Let $\langle F; +, \cdot, \exp \rangle$ be any exponential field.

Definition

$x \in F$ is **exponentially algebraic** in F iff for some $n \in \mathbb{N}$ there are:

- $\bar{x} = (x_1, \dots, x_n) \in F^n$
- $f_1, \dots, f_n \in \mathbb{Z}[\bar{X}, e^{\bar{X}}]$

such that

- $x = x_1$
- $f_i(\bar{x}, e^{\bar{x}}) = 0$ for each $i = 1, \dots, n$

- $$\begin{vmatrix} \frac{\partial f_1}{\partial X_1} & \cdots & \frac{\partial f_1}{\partial X_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial X_1} & \cdots & \frac{\partial f_n}{\partial X_n} \end{vmatrix}(\bar{x}) \neq 0$$

Exponentially Transcendental in $F \iff$ not exponentially algebraic in F

Exponential Transcendence - general definition

Let $\langle F; +, \cdot, \exp \rangle$ be any exponential field.

Definition

$x \in F$ is **exponentially algebraic** in F iff for some $n \in \mathbb{N}$ there are:

- $\bar{x} = (x_1, \dots, x_n) \in F^n$
- $f_1, \dots, f_n \in \mathbb{Z}[\bar{X}, e^{\bar{X}}]$

such that

- $x = x_1$
- $f_i(\bar{x}, e^{\bar{x}}) = 0$ for each $i = 1, \dots, n$

- $$\begin{vmatrix} \frac{\partial f_1}{\partial X_1} & \cdots & \frac{\partial f_1}{\partial X_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial X_1} & \cdots & \frac{\partial f_n}{\partial X_n} \end{vmatrix}(\bar{x}) \neq 0$$

Exponentially Transcendental in $F \iff$ not exponentially algebraic in F

A generalization

First Theorem

Let $\lambda \in \mathbb{R}$ be exponentially transcendental, let $\bar{y} \in (\mathbb{R}_{>0})^n$, and suppose \bar{y} is multiplicatively independent. Then

$$\text{td}(\mathbb{Q}(\bar{y}, \bar{y}^\lambda, \lambda)/\mathbb{Q}(\lambda)) \geq n.$$

Theorem (BKW)

F any exponential field, $\lambda \in F$ exponentially transcendental, $\bar{x} \in F^n$ such that $\exp(\bar{x})$ is multiplicatively independent. Then

$$\text{td}(\mathbb{Q}(\exp(\bar{x}), \exp(\lambda\bar{x}), \lambda)/\mathbb{Q}(\lambda)) \geq n.$$

A generalization

First Theorem

Let $\lambda \in \mathbb{R}$ be exponentially transcendental, let $\bar{y} \in (\mathbb{R}_{>0})^n$, and suppose \bar{y} is multiplicatively independent. Then

$$\text{td}(\mathbb{Q}(\bar{y}, \bar{y}^\lambda, \lambda)/\mathbb{Q}(\lambda)) \geq n.$$

Theorem (BKW)

F any exponential field, $\lambda \in F$ exponentially transcendental, $\bar{x} \in F^n$ such that $\exp(\bar{x})$ is multiplicatively independent. Then

$$\text{td}(\mathbb{Q}(\exp(\bar{x}), \exp(\lambda\bar{x}), \lambda)/\mathbb{Q}(\lambda)) \geq n.$$

Several powers

For $A \subseteq F$, can define $\text{ecl } A$, the **exponential algebraic closure** of A .

Theorem (Kirby)

*ecl is a pregeometry on any exponential field. Thus we have a notion of **independence**.*

Theorem (BKW – question of Zilber)

Let $\lambda_1, \dots, \lambda_m$ be ecl-independent in F , let $\bar{z} \in F^n$, and write \ker for the kernel of \exp .

$$\text{td}(\mathbb{Q}(\exp(\bar{z}), \bar{\lambda})/\mathbb{Q}(\bar{\lambda})) + \text{ldim}_{\mathbb{Q}(\bar{\lambda})}(\bar{z}/\ker) - \text{ldim}_{\mathbb{Q}}(\bar{z}/\ker) \geq 0$$

Several powers

For $A \subseteq F$, can define $\text{ecl } A$, the **exponential algebraic closure** of A .

Theorem (Kirby)

*ecl is a pregeometry on any exponential field. Thus we have a notion of **independence**.*

Theorem (BKW – question of Zilber)

Let $\lambda_1, \dots, \lambda_m$ be ecl-independent in F , let $\bar{z} \in F^n$, and write \ker for the kernel of \exp .

$$\text{td}(\mathbb{Q}(\exp(\bar{z}), \bar{\lambda})/\mathbb{Q}(\bar{\lambda})) + \text{ldim}_{\mathbb{Q}(\bar{\lambda})}(\bar{z}/\ker) - \text{ldim}_{\mathbb{Q}}(\bar{z}/\ker) \geq 0$$

Several powers

For $A \subseteq F$, can define $\text{ecl } A$, the **exponential algebraic closure** of A .

Theorem (Kirby)

*ecl is a pregeometry on any exponential field. Thus we have a notion of **independence**.*

Theorem (BKW – question of Zilber)

Let $\lambda_1, \dots, \lambda_m$ be ecl-independent in F , let $\bar{z} \in F^n$, and write \ker for the kernel of \exp .

$$\text{td}(\mathbb{Q}(\exp(\bar{z}), \bar{\lambda})/\mathbb{Q}(\bar{\lambda})) + \text{ldim}_{\mathbb{Q}(\bar{\lambda})}(\bar{z}/\ker) - \text{ldim}_{\mathbb{Q}}(\bar{z}/\ker) \geq 0$$

Outline

1 Motivation – Decidability

2 Schanuel Properties

3 Proofs

Outline of the proof in the real case.

Step 1

Take $\lambda \in \mathbb{R}$ exponentially transcendental, $B \cup \{\lambda\}$ an ecl-basis for \mathbb{R} , $C = \text{ecl}(B)$. Then for any $\bar{z} \in \mathbb{R}^n$,

$$\text{td}(\bar{z}, \lambda, \exp(\bar{z}), \exp(\lambda)/C) - \text{l dim}_{\mathbb{Q}}(\bar{z}, \lambda/C) \geq 1$$

- 1 For each $a \in \mathbb{R}$, there is a C -definable function $\theta : \mathbb{R} \rightarrow \mathbb{R}$ with $\theta(\lambda) = a$.
- 2 If θ, ψ are two such, o-minimality implies $\{x \in \mathbb{R} \mid \theta(x) = \psi(x)\}$ contains an interval around λ .
- 3 Similarly, θ is differentiable near λ .
- 4 Define $\partial : \mathbb{R} \rightarrow \mathbb{R}$ by $a \mapsto \frac{d\theta}{dx}(\lambda)$, where $\theta(\lambda) = a$.
- 5 ∂ is a well-defined derivation on \mathbb{R} , vanishing on C .
- 6 $\partial e^{z_i} = e^{z_i} \partial z_i$, each i , and $\partial e^\lambda = e^\lambda \partial \lambda$.
- 7 Result follows from Ax's differential field version of Schanuel's Conjecture.

Outline of the proof in the real case.

Step 1

Take $\lambda \in \mathbb{R}$ exponentially transcendental, $B \cup \{\lambda\}$ an ecl-basis for \mathbb{R} , $C = \text{ecl}(B)$. Then for any $\bar{z} \in \mathbb{R}^n$,

$$\text{td}(\bar{z}, \lambda, \exp(\bar{z}), \exp(\lambda)/C) - \text{ldim}_{\mathbb{Q}}(\bar{z}, \lambda/C) \geq 1$$

- 1 For each $a \in \mathbb{R}$, there is a C -definable function $\theta : \mathbb{R} \rightarrow \mathbb{R}$ with $\theta(\lambda) = a$.
- 2 If θ, ψ are two such, o-minimality implies $\{x \in \mathbb{R} \mid \theta(x) = \psi(x)\}$ contains an interval around λ .
- 3 Similarly, θ is differentiable near λ .
- 4 Define $\partial : \mathbb{R} \rightarrow \mathbb{R}$ by $a \mapsto \frac{d\theta}{dx}(\lambda)$, where $\theta(\lambda) = a$.
- 5 ∂ is a well-defined derivation on \mathbb{R} , vanishing on C .
- 6 $\partial e^{z_i} = e^{z_i} \partial z_i$, each i , and $\partial e^\lambda = e^\lambda \partial \lambda$.
- 7 Result follows from Ax's differential field version of Schanuel's Conjecture.

Outline of the proof in the real case.

Step 1

Take $\lambda \in \mathbb{R}$ exponentially transcendental, $B \cup \{\lambda\}$ an ecl-basis for \mathbb{R} , $C = \text{ecl}(B)$. Then for any $\bar{z} \in \mathbb{R}^n$,

$$\text{td}(\bar{z}, \lambda, \exp(\bar{z}), \exp(\lambda)/C) - \text{ldim}_{\mathbb{Q}}(\bar{z}, \lambda/C) \geq 1$$

- 1 For each $a \in \mathbb{R}$, there is a C -definable function $\theta : \mathbb{R} \rightarrow \mathbb{R}$ with $\theta(\lambda) = a$.
- 2 If θ, ψ are two such, o-minimality implies $\{x \in \mathbb{R} \mid \theta(x) = \psi(x)\}$ contains an interval around λ .
- 3 Similarly, θ is differentiable near λ .
- 4 Define $\partial : \mathbb{R} \rightarrow \mathbb{R}$ by $a \mapsto \frac{d\theta}{dx}(\lambda)$, where $\theta(\lambda) = a$.
- 5 ∂ is a well-defined derivation on \mathbb{R} , vanishing on C .
- 6 $\partial e^{z_i} = e^{z_i} \partial z_i$, each i , and $\partial e^\lambda = e^\lambda \partial \lambda$.
- 7 Result follows from Ax's differential field version of Schanuel's Conjecture.

Outline of the proof in the real case.

Step 1

Take $\lambda \in \mathbb{R}$ exponentially transcendental, $B \cup \{\lambda\}$ an ecl-basis for \mathbb{R} , $C = \text{ecl}(B)$. Then for any $\bar{z} \in \mathbb{R}^n$,

$$\text{td}(\bar{z}, \lambda, \exp(\bar{z}), \exp(\lambda)/C) - \text{ldim}_{\mathbb{Q}}(\bar{z}, \lambda/C) \geq 1$$

- 1 For each $a \in \mathbb{R}$, there is a C -definable function $\theta : \mathbb{R} \rightarrow \mathbb{R}$ with $\theta(\lambda) = a$.
- 2 If θ, ψ are two such, o-minimality implies $\{x \in \mathbb{R} \mid \theta(x) = \psi(x)\}$ contains an interval around λ .
- 3 Similarly, θ is differentiable near λ .
- 4 Define $\partial : \mathbb{R} \rightarrow \mathbb{R}$ by $a \mapsto \frac{d\theta}{dx}(\lambda)$, where $\theta(\lambda) = a$.
- 5 ∂ is a well-defined derivation on \mathbb{R} , vanishing on C .
- 6 $\partial e^{z_i} = e^{z_i} \partial z_i$, each i , and $\partial e^\lambda = e^\lambda \partial \lambda$.
- 7 Result follows from Ax's differential field version of Schanuel's Conjecture.

Outline of the proof in the real case.

Step 1

Take $\lambda \in \mathbb{R}$ exponentially transcendental, $B \cup \{\lambda\}$ an ecl-basis for \mathbb{R} , $C = \text{ecl}(B)$. Then for any $\bar{z} \in \mathbb{R}^n$,

$$\text{td}(\bar{z}, \lambda, \exp(\bar{z}), \exp(\lambda)/C) - \text{ldim}_{\mathbb{Q}}(\bar{z}, \lambda/C) \geq 1$$

- 1 For each $a \in \mathbb{R}$, there is a C -definable function $\theta : \mathbb{R} \rightarrow \mathbb{R}$ with $\theta(\lambda) = a$.
- 2 If θ, ψ are two such, o-minimality implies $\{x \in \mathbb{R} \mid \theta(x) = \psi(x)\}$ contains an interval around λ .
- 3 Similarly, θ is differentiable near λ .
- 4 Define $\partial : \mathbb{R} \rightarrow \mathbb{R}$ by $a \mapsto \frac{d\theta}{dx}(\lambda)$, where $\theta(\lambda) = a$.
- 5 ∂ is a well-defined derivation on \mathbb{R} , vanishing on C .
- 6 $\partial e^{z_i} = e^{z_i} \partial z_i$, each i , and $\partial e^\lambda = e^\lambda \partial \lambda$.
- 7 Result follows from Ax's differential field version of Schanuel's Conjecture.

Outline of the proof in the real case.

Step 1

Take $\lambda \in \mathbb{R}$ exponentially transcendental, $B \cup \{\lambda\}$ an ecl-basis for \mathbb{R} , $C = \text{ecl}(B)$. Then for any $\bar{z} \in \mathbb{R}^n$,

$$\text{td}(\bar{z}, \lambda, \exp(\bar{z}), \exp(\lambda)/C) - \text{ldim}_{\mathbb{Q}}(\bar{z}, \lambda/C) \geq 1$$

- 1 For each $a \in \mathbb{R}$, there is a C -definable function $\theta : \mathbb{R} \rightarrow \mathbb{R}$ with $\theta(\lambda) = a$.
- 2 If θ, ψ are two such, o-minimality implies $\{x \in \mathbb{R} \mid \theta(x) = \psi(x)\}$ contains an interval around λ .
- 3 Similarly, θ is differentiable near λ .
- 4 Define $\partial : \mathbb{R} \rightarrow \mathbb{R}$ by $a \mapsto \frac{d\theta}{dx}(\lambda)$, where $\theta(\lambda) = a$.
- 5 ∂ is a well-defined derivation on \mathbb{R} , vanishing on C .
- 6 $\partial e^{z_i} = e^{z_i} \partial z_i$, each i , and $\partial e^\lambda = e^\lambda \partial \lambda$.
- 7 Result follows from Ax's differential field version of Schanuel's Conjecture.

Outline of the proof in the real case.

Step 1

Take $\lambda \in \mathbb{R}$ exponentially transcendental, $B \cup \{\lambda\}$ an ecl-basis for \mathbb{R} , $C = \text{ecl}(B)$. Then for any $\bar{z} \in \mathbb{R}^n$,

$$\text{td}(\bar{z}, \lambda, \exp(\bar{z}), \exp(\lambda)/C) - \text{ldim}_{\mathbb{Q}}(\bar{z}, \lambda/C) \geq 1$$

- 1 For each $a \in \mathbb{R}$, there is a C -definable function $\theta : \mathbb{R} \rightarrow \mathbb{R}$ with $\theta(\lambda) = a$.
- 2 If θ, ψ are two such, o-minimality implies $\{x \in \mathbb{R} \mid \theta(x) = \psi(x)\}$ contains an interval around λ .
- 3 Similarly, θ is differentiable near λ .
- 4 Define $\partial : \mathbb{R} \rightarrow \mathbb{R}$ by $a \mapsto \frac{d\theta}{dx}(\lambda)$, where $\theta(\lambda) = a$.
- 5 ∂ is a well-defined derivation on \mathbb{R} , vanishing on C .
- 6 $\partial e^{z_i} = e^{z_i} \partial z_i$, each i , and $\partial e^\lambda = e^\lambda \partial \lambda$.
- 7 Result follows from Ax's differential field version of Schanuel's Conjecture.

Outline of the proof in the real case.

Step 1

Take $\lambda \in \mathbb{R}$ exponentially transcendental, $B \cup \{\lambda\}$ an ecl-basis for \mathbb{R} , $C = \text{ecl}(B)$. Then for any $\bar{z} \in \mathbb{R}^n$,

$$\text{td}(\bar{z}, \lambda, \exp(\bar{z}), \exp(\lambda)/C) - \text{ldim}_{\mathbb{Q}}(\bar{z}, \lambda/C) \geq 1$$

- 1 For each $a \in \mathbb{R}$, there is a C -definable function $\theta : \mathbb{R} \rightarrow \mathbb{R}$ with $\theta(\lambda) = a$.
- 2 If θ, ψ are two such, o-minimality implies $\{x \in \mathbb{R} \mid \theta(x) = \psi(x)\}$ contains an interval around λ .
- 3 Similarly, θ is differentiable near λ .
- 4 Define $\partial : \mathbb{R} \rightarrow \mathbb{R}$ by $a \mapsto \frac{d\theta}{dx}(\lambda)$, where $\theta(\lambda) = a$.
- 5 ∂ is a well-defined derivation on \mathbb{R} , vanishing on C .
- 6 $\partial e^{z_i} = e^{z_i} \partial z_i$, each i , and $\partial e^\lambda = e^\lambda \partial \lambda$.
- 7 Result follows from Ax's differential field version of Schanuel's Conjecture.

Step 2

$$\mathrm{td}(\exp(\bar{z})/\lambda) + \mathrm{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}) - \mathrm{ldim}_{\mathbb{Q}}(\bar{z}) \geq 0$$

Proof

We have:

$$\begin{aligned} 1 &\leq \mathrm{td}(\bar{z}, \lambda, \exp(\bar{z}), \exp(\lambda)/C) - \mathrm{ldim}_{\mathbb{Q}}(\bar{z}, \lambda/C) \\ &= \mathrm{td}(\lambda/C) + \mathrm{td}(\bar{z}/C, \lambda) + \mathrm{td}(\exp(\bar{z})/C, \lambda, \bar{z}) \\ &\quad + \mathrm{td}(\exp(\lambda)/C, \lambda, \bar{z}, \exp(\bar{z})) - \mathrm{ldim}_{\mathbb{Q}}(\lambda/C, \bar{z}) - \mathrm{ldim}_{\mathbb{Q}}(\bar{z}/C) \\ 0 &\leq \mathrm{td}(\bar{z}/C, \lambda) + \mathrm{td}(\exp(\bar{z})/\lambda) + \mathrm{td}(\exp(\lambda)/C, \exp(\bar{z})) \\ &\quad - \mathrm{ldim}_{\mathbb{Q}}(\lambda/C, \bar{z}) - \mathrm{ldim}_{\mathbb{Q}}(\bar{z}/C) \end{aligned}$$

Also and

$$\begin{aligned} \mathrm{td}(\exp(\lambda)/C, \exp(\bar{z})) &\leq \mathrm{ldim}_{\mathbb{Q}}(\lambda/C, \bar{z}) \\ \mathrm{td}(\bar{z}/C, \lambda) &\leq \mathrm{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}/C) \end{aligned}$$

Step 2

$$\mathrm{td}(\exp(\bar{z})/\lambda) + \mathrm{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}) - \mathrm{ldim}_{\mathbb{Q}}(\bar{z}) \geq 0$$

Proof

We have:

$$\begin{aligned} 1 &\leq \mathrm{td}(\bar{z}, \lambda, \exp(\bar{z}), \exp(\lambda)/C) - \mathrm{ldim}_{\mathbb{Q}}(\bar{z}, \lambda/C) \\ &= \mathrm{td}(\lambda/C) + \mathrm{td}(\bar{z}/C, \lambda) + \mathrm{td}(\exp(\bar{z})/C, \lambda, \bar{z}) \\ &\quad + \mathrm{td}(\exp(\lambda)/C, \lambda, \bar{z}, \exp(\bar{z})) - \mathrm{ldim}_{\mathbb{Q}}(\lambda/C, \bar{z}) - \mathrm{ldim}_{\mathbb{Q}}(\bar{z}/C) \\ 0 &\leq \mathrm{td}(\bar{z}/C, \lambda) + \mathrm{td}(\exp(\bar{z})/\lambda) + \mathrm{td}(\exp(\lambda)/C, \exp(\bar{z})) \\ &\quad - \mathrm{ldim}_{\mathbb{Q}}(\lambda/C, \bar{z}) - \mathrm{ldim}_{\mathbb{Q}}(\bar{z}/C) \end{aligned}$$

Also and

$$\begin{aligned} \mathrm{td}(\exp(\lambda)/C, \exp(\bar{z})) &\leq \mathrm{ldim}_{\mathbb{Q}}(\lambda/C, \bar{z}) \\ \mathrm{td}(\bar{z}/C, \lambda) &\leq \mathrm{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}/C) \end{aligned}$$

Step 2

$$\text{td}(\exp(\bar{z})/\lambda) + \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}) - \text{ldim}_{\mathbb{Q}}(\bar{z}) \geq 0$$

Proof continued

Putting these together we get

$$\text{td}(\exp(\bar{z})/\lambda) + \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}/C) - \text{ldim}_{\mathbb{Q}}(\bar{z}/C) \geq 0$$

But $\mathbb{Q}(\lambda)$ is linearly disjoint from C over \mathbb{Q} , so

$$\text{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}/C) - \text{ldim}_{\mathbb{Q}}(\bar{z}/C) \leq \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}) - \text{ldim}_{\mathbb{Q}}(\bar{z})$$

which gives the result. □

Step 3

A similar argument shows for any \bar{x} :

$$\text{ldim}_{\mathbb{Q}}(\lambda\bar{x}/\bar{x}) \geq \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{x})$$

Step 2

$$\text{td}(\exp(\bar{z})/\lambda) + \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}) - \text{ldim}_{\mathbb{Q}}(\bar{z}) \geq 0$$

Proof continued

Putting these together we get

$$\text{td}(\exp(\bar{z})/\lambda) + \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}/C) - \text{ldim}_{\mathbb{Q}}(\bar{z}/C) \geq 0$$

But $\mathbb{Q}(\lambda)$ is linearly disjoint from C over \mathbb{Q} , so

$$\text{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}/C) - \text{ldim}_{\mathbb{Q}}(\bar{z}/C) \leq \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}) - \text{ldim}_{\mathbb{Q}}(\bar{z})$$

which gives the result. □

Step 3

A similar argument shows for any \bar{x} :

$$\text{ldim}_{\mathbb{Q}}(\lambda\bar{x}/\bar{x}) \geq \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{x})$$

Step 2

$$\text{td}(\exp(\bar{z})/\lambda) + \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}) - \text{ldim}_{\mathbb{Q}}(\bar{z}) \geq 0$$

Proof continued

Putting these together we get

$$\text{td}(\exp(\bar{z})/\lambda) + \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}/C) - \text{ldim}_{\mathbb{Q}}(\bar{z}/C) \geq 0$$

But $\mathbb{Q}(\lambda)$ is linearly disjoint from C over \mathbb{Q} , so

$$\text{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}/C) - \text{ldim}_{\mathbb{Q}}(\bar{z}/C) \leq \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}) - \text{ldim}_{\mathbb{Q}}(\bar{z})$$

which gives the result. □

Step 3

A similar argument shows for any \bar{x} :

$$\text{ldim}_{\mathbb{Q}}(\lambda\bar{x}/\bar{x}) \geq \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{x})$$

Theorem

For $\bar{y} \in \mathbb{R}_{>0}^n$ multiplicatively independent,

$$\text{td}(\bar{y}, \bar{y}^\lambda / \lambda) \geq n$$

Proof.

- Step 2: $\forall \bar{z} \quad \text{td}(\exp(\bar{z})/\lambda) + \text{l dim}_{\mathbb{Q}(\lambda)}(\bar{z}) - \text{l dim}_{\mathbb{Q}}(\bar{z}) \geq 0$
- Take $\bar{x} = \log \bar{y} \quad \bar{z} = (\bar{x}, \lambda \bar{x})$
- Then

$$\begin{aligned} \text{td}(\bar{y}, \bar{y}^\lambda / \lambda) &\geq \text{l dim}_{\mathbb{Q}}(\bar{x}, \lambda \bar{x}) - \text{l dim}_{\mathbb{Q}(\lambda)}(\bar{x}, \lambda \bar{x}) \\ &\geq \text{l dim}_{\mathbb{Q}}(\bar{x}) + \text{l dim}_{\mathbb{Q}}(\lambda \bar{x} / \bar{x}) - \text{l dim}_{\mathbb{Q}(\lambda)}(\bar{x}) \\ &\geq n \end{aligned}$$

as \bar{x} is \mathbb{Q} -linearly independent and by step 3.



Theorem

For $\bar{y} \in \mathbb{R}_{>0}^n$ multiplicatively independent,

$$\text{td}(\bar{y}, \bar{y}^\lambda / \lambda) \geq n$$

Proof.

- Step 2: $\forall \bar{z} \quad \text{td}(\exp(\bar{z})/\lambda) + \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}) - \text{ldim}_{\mathbb{Q}}(\bar{z}) \geq 0$
- Take $\bar{x} = \log \bar{y} \quad \bar{z} = (\bar{x}, \lambda \bar{x})$
- Then

$$\begin{aligned} \text{td}(\bar{y}, \bar{y}^\lambda / \lambda) &\geq \text{ldim}_{\mathbb{Q}}(\bar{x}, \lambda \bar{x}) - \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{x}, \lambda \bar{x}) \\ &\geq \text{ldim}_{\mathbb{Q}}(\bar{x}) + \text{ldim}_{\mathbb{Q}}(\lambda \bar{x} / \bar{x}) - \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{x}) \\ &\geq n \end{aligned}$$

as \bar{x} is \mathbb{Q} -linearly independent and by step 3.



Theorem

For $\bar{y} \in \mathbb{R}_{>0}^n$ multiplicatively independent,

$$\text{td}(\bar{y}, \bar{y}^\lambda / \lambda) \geq n$$

Proof.

- Step 2: $\forall \bar{z} \quad \text{td}(\exp(\bar{z})/\lambda) + \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}) - \text{ldim}_{\mathbb{Q}}(\bar{z}) \geq 0$
- Take $\bar{x} = \log \bar{y} \quad \bar{z} = (\bar{x}, \lambda \bar{x})$
- Then

$$\begin{aligned} \text{td}(\bar{y}, \bar{y}^\lambda / \lambda) &\geq \text{ldim}_{\mathbb{Q}}(\bar{x}, \lambda \bar{x}) - \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{x}, \lambda \bar{x}) \\ &\geq \text{ldim}_{\mathbb{Q}}(\bar{x}) + \text{ldim}_{\mathbb{Q}}(\lambda \bar{x} / \bar{x}) - \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{x}) \\ &\geq n \end{aligned}$$

as \bar{x} is \mathbb{Q} -linearly independent and by step 3.



Theorem

For $\bar{y} \in \mathbb{R}_{>0}^n$ multiplicatively independent,

$$\text{td}(\bar{y}, \bar{y}^\lambda / \lambda) \geq n$$

Proof.

- Step 2: $\forall \bar{z} \quad \text{td}(\exp(\bar{z})/\lambda) + \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}) - \text{ldim}_{\mathbb{Q}}(\bar{z}) \geq 0$
- Take $\bar{x} = \log \bar{y} \quad \bar{z} = (\bar{x}, \lambda \bar{x})$
- Then

$$\begin{aligned} \text{td}(\bar{y}, \bar{y}^\lambda / \lambda) &\geq \text{ldim}_{\mathbb{Q}}(\bar{x}, \lambda \bar{x}) - \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{x}, \lambda \bar{x}) \\ &\geq \text{ldim}_{\mathbb{Q}}(\bar{x}) + \text{ldim}_{\mathbb{Q}}(\lambda \bar{x} / \bar{x}) - \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{x}) \\ &\geq n \end{aligned}$$

as \bar{x} is \mathbb{Q} -linearly independent and by step 3.



Theorem

For $\bar{y} \in \mathbb{R}_{>0}^n$ multiplicatively independent,

$$\text{td}(\bar{y}, \bar{y}^\lambda / \lambda) \geq n$$

Proof.

- Step 2: $\forall \bar{z} \quad \text{td}(\exp(\bar{z})/\lambda) + \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{z}) - \text{ldim}_{\mathbb{Q}}(\bar{z}) \geq 0$
- Take $\bar{x} = \log \bar{y} \quad \bar{z} = (\bar{x}, \lambda \bar{x})$
- Then

$$\begin{aligned} \text{td}(\bar{y}, \bar{y}^\lambda / \lambda) &\geq \text{ldim}_{\mathbb{Q}}(\bar{x}, \lambda \bar{x}) - \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{x}, \lambda \bar{x}) \\ &\geq \text{ldim}_{\mathbb{Q}}(\bar{x}) + \text{ldim}_{\mathbb{Q}}(\lambda \bar{x} / \bar{x}) - \text{ldim}_{\mathbb{Q}(\lambda)}(\bar{x}) \\ &\geq n \end{aligned}$$

as \bar{x} is \mathbb{Q} -linearly independent and by step 3.

